



Exteris™ Stressgard®

Product

information sheet

// BEST USES

Exteris™ Stressgard® is formulated with a novel SDHI active ingredient, fluopyram, that is combined with trifloxystrobin to provide long-lasting preventative and curative control of foliar diseases including dollar spot, anthracnose, brown patch, and leaf spot. The unique formulation delivers enhanced plant health benefits along with improved turf quality. The flexibility of Exteris Stressgard allows for applications throughout the year and under a wide variety of environmental conditions.

// KEY STRENGTHS

Preventative and fast curative dollar spot control on fairways from 21-28 days.

Attributes of Exteris Stressgard

- Broad-spectrum disease control – dollar spot, anthracnose, brown patch, leaf spot.
- Program flexibility for greens, tees and fairways.
- Enhanced turf quality and plant health benefits associated with Stressgard® Formulation Technology.
- Quick dry time and absorption.
- Fast curative activity across range of spray volumes.
- Dew mitigation properties.

Solutions for Tough Diseases

- Preventative and fast curative activity.
- Controls DMI resistant dollar spot.
- Can be used for summer dollar spot, anthracnose and brown patch control without negative growth regulating effects.
- Promotes plant health throughout the summer disease season.

// HOW TO USE EXTERIS STRESSGARD

Apply Exteris Stressgard when conditions are favourable for disease as part of an integrated management program on greens, tees and fairways. Exteris Stressgard can be combined with Signature XTRA™ Stressgard® in an anthracnose management program for enhanced turf quality.

NOTE: Exteris Stressgard is compatible with most fungicide tank-mixes, but a jar test is always recommended. Exteris Stressgard should NOT be mixed with Daconil® 2787 due to incompatibility concerns.

// Active Ingredients

fluopyram
(12.5 g/L) + trifloxystrobin
(12.5 g/L)

// FRAC Code

7 + 11 (SDHI + QoI)

// Systemicity

acropetal

// Formulation

suspension
concentrate

// Signal Word

caution


Exteris Stressgard Rate Chart

Use Rate (g/100 m ²)	Coverage Per Container (100 m ²)	Coverage Per Container (Acres)	Grams Trifloxystrobin (Per Acre)	Grams Fluopyram (Per Acre)	Applications Allowed at Rate (Per Year)
140	71	1.75	708	708	2.8
170	58.8	1.45	859.9	859.9	2.4
200	50	1.24	1011.6	1011.6	2


Key Diseases Controlled

Key Diseases Controlled	Rates mL/100 m ²	Application Interval (Days)
Dollar spot	140-200	14-21 (greens and tees) 21-28 (fairways)
Brown patch	140-200	14-21
Leaf spot (<i>Bipolaris</i> spp., <i>Drechslera</i> spp.)	140-200	14-21
Anthracnose (foliar blight)*	140	14-21
Anthracnose (foliar blight and basal rot)	200	14-21

See Exteris Stressgard label for complete reference on control options and applications instructions. Always read and carefully follow label instructions.



Control
Photo: Dr. Paul Giordano - Bayer



Exteris Stressgard, 200 mL, 28 days
Photo: Dr. Paul Giordano - Bayer



Western Canada:
Josey Groenveld
403.463.2742

Ontario:
Darcy Olds
905.319.8981

Ontario:
Keith Bartlett
226.821.2356

Atlantic/Quebec/
Eastern Ontario:
Jonathan Albert
226.821.0077

Quebec:
Normand Drapeau
514.949.2467

Green Solutions Specialist:
Jesse Benelli Ph.D.
905.302.0676

www.bayeres.ca 1-888-283-6847 @BayerGolfCA